




## HPU2 Journal of Sciences: Natural Sciences and Technology

Journal homepage: <https://sj.hpu2.edu.vn>



*Article type: Research article*

### On the second-order sufficient optimality condition in nonconvex multiobjective optimization problems

Van-Tuyen Nguyen<sup>a\*</sup>, Thi-Yen Nguyen<sup>b</sup>

<sup>a</sup>Hanoi Pedagogical University 2, Vinh Phuc, Vietnam

<sup>b</sup>Phenikaa University, Hanoi, Vietnam

#### Abstract

The study of second-order optimality conditions is one of the most important topics in optimization theory and attracting the attention and interest of many authors. In this paper, we introduce a novel solution concept called “essential local efficient solutions of second-order” for nonconvex constrained multiobjective optimization problems. We then show that the new solution concept is stronger than the quadratic growth condition and under a mild constraint qualification, these solution concepts are equivalent. By using the second subderivative, we derive a sufficient optimality condition for a feasible solution to become an essential local efficient solution of second-order for the considered problem. Examples are provided to illustrate the obtained results.

**Keywords:** Essential local efficient solutions of second-order, second subderivative, second-order sufficient optimality condition

#### 1. Introduction

Second-order optimality conditions have long been recognized as an important tool in optimization theory and, in recent years, have been developed rapidly, see, for example [1]–[16]. It is well known that first-order optimality conditions are usually not sufficient for optimality except in the case of convex optimization problems. Second-order optimality conditions not only complement first-order ones in eliminating non-optimal solutions, but they also provide criteria for recognizing the optimality at a given feasible solution.

\* Corresponding author, E-mail: [nguyenvantuyen83@hpu2.edu.vn](mailto:nguyenvantuyen83@hpu2.edu.vn)

<https://doi.org/10.56764/hpu2.jos.2024.3.3.60-69>

Received date: 23-9-2024 ; Revised date: 28-10-2024 ; Accepted date: 08-11-2024

This is licensed under the CC BY-NC 4.0

In this paper, we will focus on second-order sufficient optimality conditions for the following constrained multiobjective optimization problem

$$\text{Min}_{\mathbb{R}_+^m} \{f(x) : g(x) \in C\}, \tag{MP}$$

where  $f : \mathbb{R}^n \rightarrow \mathbb{R}^m$  and  $g : \mathbb{R}^n \rightarrow \mathbb{R}^p$  are twice continuously differentiable mappings, and  $C \subset \mathbb{R}^p$  is a nonempty and closed set. When  $m = 1$ , the above problem is called a mathematical program problem and is denoted by (P).

The study of second-order optimality conditions for (P), when  $C$  is convex, has been completely developed by Bonnans and Shapiro [1], Cominetti [2], Rockafellar and Wets [12], Mohammadi *et al.* [10], etc. More precisely, if  $C$  is convex polyhedral, second-order optimality conditions can be expressed in term of second derivative of the Lagrangian, see, for example [1], [12]. If  $C$  lacks the polyhedrality, then an additional term is needed to capture the curvature of  $C$  and there are various tools that can be utilized for such purpose, see [2], [10].

Recently, several important problem classes which can be reformulated in the form of problem (P) with non-convex  $C$ , such as, the mathematical program with complementarity constraints, the mathematical program with semi-definite cone complementarity constraints, etc. have attracted significant attention from the optimization community, see [17]–[20]. In these papers, the authors use the so-called lower generalized support function and the second subderivative to derive necessary and sufficient optimality conditions for (P) with  $C$  nonconvex. However, to the best of our knowledge, no papers have yet investigated second-order optimality conditions for multiobjective optimization problems of the form (MP). Motivated by the works reported in [11], [17], [18], in this paper, we introduce a new solution concept called “essential local efficient solutions of second-order” for the problem (MP) and study the sufficient optimality condition for the proposed solution.

We organize the paper as follows. Section 2 contains the preliminaries and auxiliary results. In Section 3, we present a second-order sufficient optimality condition for a feasible solution to be an essential local efficient solution of second-order to (MP). Section 4 provides some concluding remarks.

## 2. Preliminaries

Throughout this work we deal with the Euclidean space  $\mathbb{R}^n$  equipped with the usual scalar product  $\langle \cdot, \cdot \rangle$  and the corresponding norm  $\|\cdot\|$ . We denote by  $B_r(x)$  the open ball centered at  $x$  with radius  $r$ . The set of all positive integer numbers is denoted by  $\mathbb{N}$ . Let  $\Omega$  be a nonempty subset in  $\mathbb{R}^n$ . The *closure*, *interior*, *convex hull*, and *conic hull* of  $\Omega$  are denoted, respectively, by  $\text{cl } \Omega$ ,  $\text{int } \Omega$ ,  $\text{conv } \Omega$ , and  $\text{cone } \Omega$ . The distance  $\text{dist}(x, \Omega)$  from a point  $x \in \mathbb{R}^n$  to  $\Omega$  is defined by

$$\text{dist}(x, \Omega) := \inf \{ \|y - x\| : y \in \Omega \} \quad \forall x \in \mathbb{R}^n.$$

The indicator function  $\delta_\Omega$  and the support function  $\sigma_\Omega$  of  $\Omega$  are defined, respectively, by

$$\begin{aligned} \sigma_\Omega(z^*) &= \sup \{ \langle z^*, z \rangle : z \in \Omega \}, \\ \delta_\Omega(x) &= \begin{cases} 0 & \text{if } x \in \Omega, \\ \infty & \text{otherwise.} \end{cases} \end{aligned}$$

**Definition 2.1.** Let  $\Omega \subset \mathbb{R}^n$ ,  $z \in \Omega$ , and  $u \in \mathbb{R}^n$ .

(i) The set *tangent/contingent cone* to  $\Omega$  at  $z$  is defined by

$$T_{\Omega}(z) := \left\{ u \in \mathbb{R}^n : \exists t_k \downarrow 0, u_k \rightarrow u \text{ with } z + t_k u_k \in \Omega \ \forall k \in \mathbb{N} \right\}.$$

(ii) The *second-order tangent set* to  $\Omega$  at  $z$  with respect to the direction  $u$  is defined by

$$T_{\Omega}^2(z, u) := \left\{ v \in \mathbb{R}^n : \exists t_k \downarrow 0, \exists v^k \rightarrow v, z + t_k u + \frac{1}{2} t_k^2 v^k \in \Omega, \ \forall k \in \mathbb{N} \right\}.$$

**Remark 2.2.** It is well-known that  $T_{\Omega}(z)$  is a nonempty closed cone. For each  $u \in \mathbb{R}^n$ , the set  $T_{\Omega}^2(z, u)$  is closed and  $T_{\Omega}^2(z, u) = \emptyset$  if  $u \notin T_{\Omega}(z)$ . However, we see that the set  $T_{\Omega}^2(z, 0) = T_{\Omega}(z)$  is always nonempty. If  $\Omega$  is convex, then we have

$$T_{\Omega}(z) = \text{cl} \left\{ d : d = \beta(x - z), \ x \in \Omega, \beta \geq 0 \right\},$$

and for each  $u \in T_{\Omega}(z)$  one has

$$T_{\Omega}^2(z, u) \subset \text{cl cone}[\text{cone}(\Omega - z) - u].$$

Moreover, if  $\Omega$  is a polyhedral convex set, then we have

$$T_{\Omega}^2(z, u) = T_{T_{\Omega}(z)}(u).$$

**Definition 2.3.** Let  $w \in \mathbb{R}^n$ . For  $\delta, \rho > 0$ ,

$$V_{\delta, \rho}(w) := \left\{ w' \in B_{\delta}(0) : \left| \|w\| w' - \|w'\| w \right| \leq \rho \|w'\| \|w\| \right\}$$

is called a *directional neighborhood of direction*  $w$ .

**Definition 2.4.** Let  $\Omega \subset \mathbb{R}^n$ ,  $z \in \Omega$ , and  $w \in T_{\Omega}(z)$ . The *proximal prenormal cone*  $\widehat{N}_{\Omega}^p(z, w)$  and the *proximal normal cone*  $\widehat{N}_{\Omega}^p(z, w)$  to  $\Omega$  at  $z$  in the direction  $w$  are defined, respectively, by

$$\widehat{N}_{\Omega}^p(z, w) := \left\{ z^* \in \mathbb{R}^n : \exists \delta, \rho, \gamma > 0 \text{ such that } \langle z^*, z' - z \rangle \leq \gamma \|z' - z\|^2 \ \forall z' \in \Omega \cap (z + V_{\delta, \rho}(w)) \right\},$$

$$\widehat{N}_{\Omega}^p(z, w) := \widehat{N}_{\Omega}^p(z, w) \cap w^{\perp}.$$

If  $w \notin T_{\Omega}(z)$ , we define  $\widehat{N}_{\Omega}^p(z, w) = \widehat{N}_{\Omega}^p(z, w) = \emptyset$ .

**Definition 2.5.** Let  $\varphi: \mathbb{R}^n \rightarrow \overline{\mathbb{R}} := [-\infty, \infty]$  be an extended real-valued function and  $z \in \mathbb{R}^n$  such that  $|\varphi(z)| < \infty$  and  $z^* \in \mathbb{R}^n$ .

(i) The *subderivative* of  $\varphi$  at  $z$  is defined by

$$d\varphi(z)(w) := \liminf_{\substack{t \downarrow 0, \\ w' \rightarrow w}} \frac{\varphi(z + tw') - \varphi(z)}{t} \quad \forall w \in \mathbb{R}^n.$$

(ii) The *second subderivative* of  $\varphi$  at  $z$  for  $z^*$  is defined by

$$d^2\varphi(z, z^*)(w) := \liminf_{\substack{t \downarrow 0, \\ w' \rightarrow w}} \frac{\varphi(z + tw') - \varphi(z) - t \langle z^*, w' \rangle}{\frac{1}{2} t^2} \quad \forall w \in \mathbb{R}^n.$$

**Remark 2.6.** (see [12]) (i) The second subderivative has the homogeneity property, i.e.,

$$d^2(\alpha\varphi)(z, z^*)(w) = \alpha d^2\varphi\left(z, \frac{z^*}{\alpha}\right)(w) \quad \forall \alpha > 0,$$

and

$$d\varphi(z)(w) > \langle z^*, w \rangle \Rightarrow d^2\varphi(z, z^*)(w) = \infty,$$

$$d\varphi(z)(w) < \langle z^*, w \rangle \Rightarrow d^2\varphi(z, z^*)(w) = -\infty.$$

(ii) If  $\varphi$  is twice differentiable at  $z$  and  $z^* = \nabla\varphi(z)$ , then we have

$$d^2\varphi(z, \nabla\varphi(z))(w) = w^T \nabla^2\varphi(z) w \quad \forall w \in \mathbb{R}^n.$$

(iii) Let  $\Omega \subset \mathbb{R}^n$ ,  $z \in \Omega$ , and  $z^* \in \mathbb{R}^n$ . Then, by definition, we have

$$d^2\delta_\Omega(z; z^*)(w) = \liminf_{\substack{t \downarrow 0 \\ w' \rightarrow w}} \frac{\delta_\Omega(z + tw') - \delta_\Omega(z) - t\langle z^*, w' \rangle}{\frac{1}{2}t^2} = \liminf_{\substack{t \downarrow 0, w' \rightarrow w \\ z + tw' \in \Omega}} \frac{-2\langle z^*, w' \rangle}{t}.$$

We now summarize some properties of the second subderivative that will be used in the next section.

**Lemma 2.7.** (see [17, Lemma 2.7]) *Let  $\varphi: \mathbb{R}^n \rightarrow \overline{\mathbb{R}}$  be a lower semicontinuous function and  $z \in \mathbb{R}^n$  such that  $|\varphi(z)| < \infty$  and  $z^* \in \mathbb{R}^n$ . Then there exist sequences  $t_k \downarrow 0$  and  $w_k \rightarrow w$  such that*

$$d\varphi(z)(w) = \lim_{k \rightarrow \infty} \frac{\varphi(z + t_k w_k) - \varphi(z)}{t_k},$$

$$d^2\varphi(z, z^*)(w) = \lim_{k \rightarrow \infty} \frac{\varphi(z + t_k w_k) - \varphi(z) - t_k \langle z^*, w_k \rangle}{\frac{1}{2}t_k^2}.$$

**Lemma 2.8.** (see [18, Proposition 2.18]) *Let  $\Omega \subset \mathbb{R}^n$ ,  $z \in \Omega$ , and  $z^*, w \in \mathbb{R}^n$ . The following statements hold:*

- (i) *If  $w \in T_\Omega(z)$  or  $\langle z^*, w \rangle < 0$ , then  $d^2\delta_\Omega(z; z^*)(w) = \infty$ .*
- (ii) *For  $w \in T_\Omega(z)$ ,  $d^2\delta_\Omega(z; z^*)(w) > -\infty$  iff  $z^* \in \widehat{N}_\Omega^p(z, w)$ .*
- (iii) *If  $d^2\delta_\Omega(z; z^*)(w)$  is finite, then  $z^* \in \widehat{N}_\Omega^p(z, w)$ .*
- (iv)  *$d^2\delta_\Omega(z; z^*)(w) \leq -\sigma_{T_\Omega^2(z, w)}(z^*)$  iff  $w \in T_\Omega(z)$  and  $\langle z^*, w \rangle \geq 0$  or  $T_\Omega^2(z, w) = \emptyset$ .*

### 3. Second-order sufficient optimality conditions

Consider the following constrained optimization problem

$$\text{Min}_{\mathbb{R}_+^m} \{f(x) : g(x) \in C\}, \tag{MP}$$

where  $f: \mathbb{R}^n \rightarrow \mathbb{R}^m$  and  $g: \mathbb{R}^n \rightarrow \mathbb{R}^p$  are twice continuously differentiable mappings with  $f(x) := (f_1(x), \dots, f_m(x))$ ,  $g(x) := (g_1(x), \dots, g_p(x))$ , and  $C \subset \mathbb{R}^n$  is a nonempty and closed set.

The feasible set of (MP) is denoted by

$$S := g^{-1}(C) = \{x \in \mathbb{R}^n : g(x) \in C\}.$$

We always assume that  $S$  is nonempty. We say that  $u$  is a *critical direction* of problem (MP) at  $\bar{x} \in S$  if

$$\begin{cases} \langle \nabla g(\bar{x}), u \rangle \in T_c(g(\bar{x})), \\ \langle \nabla f_i(\bar{x}), u \rangle \leq 0, \quad \forall i \in I := \{1, \dots, m\}. \end{cases}$$

The set of all critical directions of (MP) at  $\bar{x} \in S$  is denoted by  $K(\bar{x})$ . We say that the set-valued mapping  $x \mapsto g(x) - C$  is *metrically subregular* at  $(\bar{x}, 0)$  in direction  $u \in \mathbb{R}^n$  if there exist  $\kappa, \delta, \rho > 0$  such that

$$\text{dist}(x, S) \leq \kappa \text{dist}(g(x), C) \quad \forall x \in \bar{x} + V_{\delta, \rho}(u).$$

The *generalized Lagrangian*  $L : \mathbb{R}^n \times \mathbb{R}_+^m \times \mathbb{R}^p \rightarrow \mathbb{R}$  with respect to the problem (MP) is given as

$$L(x, \lambda, \mu) := \langle \lambda, f(x) \rangle + \langle \mu, g(x) \rangle.$$

We now introduce the concept of essential local efficient solutions of second-order for (MOP) inspired by the work of Penot [11].

**Definition 3.1.** Let  $\bar{x} \in S$ . We say that:

(i)  $\bar{x}$  is a *local efficient solution* of (MP) if there exists  $\delta > 0$  such that there is no  $x \in S \cap B_\delta(\bar{x})$  satisfying

$$f(x) \in f(\bar{x}) - \mathbb{R}_+^m \setminus \{0\}.$$

(ii)  $\bar{x}$  satisfies the *quadratic growth condition* if there exist two positive numbers  $\beta > 0$  and  $\delta > 0$  such that

$$\psi(x) := \max\{f_1(x) - f_1(\bar{x}), \dots, f_m(x) - f_m(\bar{x})\} \geq \beta \|x - \bar{x}\|^2 \quad \forall x \in S \cap B_\delta(\bar{x}).$$

(iii)  $\bar{x}$  is an *essential local efficient solution of second-order* for problem (MP) if there exist two positive numbers  $\gamma > 0$  and  $\delta > 0$  such that

$$\varphi(x) := \max\{f_1(x) - f_1(\bar{x}), \dots, f_m(x) - f_m(\bar{x}), \text{dist}(g(x), C)\} \geq \gamma \|x - \bar{x}\|^2 \quad \forall x \in B_\delta(\bar{x}).$$

The following result gives the relationships between above solution concepts.

**Proposition 3.2.** Consider the following statements:

(i)  $\bar{x}$  is a local efficient solution of (MP).

(ii)  $\bar{x}$  satisfies the quadratic growth condition.

(iii)  $\bar{x}$  is an essential local efficient solution of second-order for problem (MP).

Then the implications (iii)  $\Rightarrow$  (ii) and (ii)  $\Rightarrow$  (i) always hold. Furthermore, if the mapping  $x \mapsto g(x) - C$  is metrically subregular at  $(\bar{x}, 0)$  in every critical direction  $u \in K(\bar{x}) \setminus \{0\}$ , then the implication (ii)  $\Rightarrow$  (iii) is also valid.

*Proof.* (iii)  $\Rightarrow$  (ii): The proof follows immediately from the definitions.

(ii)  $\Rightarrow$  (i): Suppose, for the sake of contradiction that  $\bar{x}$  satisfies the quadratic growth condition but not is a local efficient solution of (MP). Then, by definition, there exist two positive numbers

$\beta > 0$ ,  $\delta > 0$ , and some  $x_0 \in S \cap B_\delta(\bar{x})$  such that

$$\max\{f_1(x) - f_1(\bar{x}), \dots, f_m(x) - f_m(\bar{x})\} \geq \beta \|x - \bar{x}\|^2 \quad \forall x \in S \cap B_\delta(\bar{x}). \quad (1)$$

and

$$f_i(x_0) \leq f_i(\bar{x}) \quad \forall i \in I \quad (2)$$

with at least one strict inequality. By (2),  $x_0 \neq \bar{x}$ . This and (1) imply that

$$\max\{f_1(x_0) - f_1(\bar{x}), \dots, f_m(x_0) - f_m(\bar{x})\} \geq \gamma \|x_0 - \bar{x}\|^2 > 0,$$

contrary to (2).

We now show that (ii)  $\Rightarrow$  (iii) under the metric subregularity of the mapping  $g(\cdot) - C$  at  $(\bar{x}, 0)$ . Suppose, for the sake of contradiction that  $\bar{x}$  is not an essential local efficient solution of second-order for (P). Then for any  $k \in \mathbb{N}$ , there exists  $x_k \in B_{1/k}(\bar{x})$  such that

$$\varphi(x_k) = \max\{f_1(x_k) - f_1(\bar{x}), \dots, f_m(x_k) - f_m(\bar{x}), \text{dist}(g(x_k), C)\} < \frac{1}{k} \|x_k - \bar{x}\|^2. \quad (3)$$

It is clear that  $x_k \neq \bar{x}$  for all  $k \in \mathbb{N}$  and  $x_k \rightarrow \bar{x}$ . Hence,

$$\liminf_{k \rightarrow \infty} \frac{f_i(x_k) - f_i(\bar{x})}{\|x_k - \bar{x}\|^2} \leq 0 \quad \forall i \in I. \quad (4)$$

By passing to a subsequence we may assume that  $\frac{x_k - \bar{x}}{\|x_k - \bar{x}\|} \rightarrow u \neq 0$  as  $k \rightarrow \infty$ . We first claim that  $u \in K(\bar{x})$ . Indeed, for each  $k \in \mathbb{N}$ , put  $t_k := \|x_k - \bar{x}\|$  and  $u_k := \frac{1}{t_k}(x_k - \bar{x})$ . Then it follows from (3) that  $\text{dist}(g(x_k), C) < \frac{1}{k} t_k^2$ . Hence, there exists  $y_k \in C$  such that  $\|g(x_k) - y_k\| < \frac{1}{k} t_k^2$ . Put  $r_k := \frac{y_k - g(x_k)}{t_k^2}$ . Then we see that  $\|r_k\| \rightarrow 0$  as  $k \rightarrow \infty$  and  $y_k = g(x_k) + t_k^2 r_k \in C$  for all  $k \in \mathbb{N}$ . Since  $x_k = \bar{x} + t_k u_k$ , by Taylor's expansion, we have

$$g(x_k) = g(\bar{x}) + t_k \langle \nabla g(\bar{x}), u_k \rangle + o(t_k).$$

Hence,

$$v_k := \frac{g(x_k) + t_k^2 r_k - g(\bar{x})}{t_k} = \frac{t_k \langle \nabla g(\bar{x}), u_k \rangle + o(t_k) + t_k^2 r_k}{t_k} = \langle \nabla g(\bar{x}), u_k \rangle + t_k r_k + \frac{o(t_k)}{t_k} \rightarrow \langle \nabla g(\bar{x}), u \rangle$$

as  $k \rightarrow \infty$ . Furthermore,  $g(\bar{x}) + t_k v_k = g(x_k) + t_k^2 r_k = y_k \in C$  for all  $k \in \mathbb{N}$ . Hence,  $\langle \nabla g(\bar{x}), u \rangle \in T_C(g(\bar{x}))$ . We deduce again from (3) that  $f_i(x_k) - f_i(\bar{x}) < \frac{1}{k} t_k^2$  for all  $k \in \mathbb{N}$  and  $i \in I$ . This and the Taylor's expansion of  $f_i$  at  $\bar{x}$ ,  $i \in I$ , imply that  $\langle \nabla f_i(\bar{x}), u \rangle \leq 0, \forall i \in I$ , and hence,  $u \in K(\bar{x})$ , as required.

Now, by the assumption on the metric subregularity of the mapping  $g(\cdot) - C$  at  $(\bar{x}, 0)$  in the

direction  $u \in K(\bar{x})$ , there exists  $\ell > 0$  such that for all  $k$  large enough we can find some  $\hat{x}_k \in S$  with

$$\|\hat{x}_k - x_k\| \leq \ell \operatorname{dist}(g(x_k), C) < \frac{\ell}{k} t_k^2.$$

Clearly,  $\hat{x}_k \rightarrow \bar{x}$  as  $k \rightarrow \infty$  and  $\|\hat{x}_k - x_k\| \leq o(\|x_k - \bar{x}\|^2)$ . Since  $\bar{x}$  satisfies the quadratic growth condition, there exist two positive numbers  $\beta > 0$  and  $\delta > 0$  satisfying (1). Hence, for each  $k$  large enough, there exists  $i_k \in I$  such that

$$f_{i_k}(\hat{x}_k) - f_{i_k}(\bar{x}) \geq \beta \|\hat{x}_k - \bar{x}\|^2. \tag{5}$$

Let  $I_k$  be the set of all indices  $i_k \in I$  satisfying (5). Since  $I_k \subset I$  for all  $k$ , without any loss of generality, we may assume that  $I_k = \bar{I}$  is constant for all  $k \in \mathbb{N}$  large enough. Fix  $i \in \bar{I}$ , then one has

$$f_i(\hat{x}_k) - f_i(\bar{x}) \geq \beta \|\hat{x}_k - \bar{x}\|^2$$

for all  $k$  large enough. Clearly,  $f_i$  is locally Lipschitz around  $\bar{x}$  with some constant  $l_i > 0$ . Hence,

$$\begin{aligned} 0 < \beta &\leq \liminf_{k \rightarrow \infty} \frac{f_i(\hat{x}_k) - f_i(\bar{x})}{\|\hat{x}_k - \bar{x}\|^2} \leq \liminf_{k \rightarrow \infty} \frac{f_i(x_k) - f_i(\bar{x}) + l_i \|\hat{x}_k - x_k\|}{(\|x_k - \bar{x}\| - \|\hat{x}_k - x_k\|)^2} \\ &= \liminf_{k \rightarrow \infty} \frac{f_i(x_k) - f_i(\bar{x}) + o(\|\hat{x}_k - x_k\|^2)}{\|\hat{x}_k - x_k\|^2 + o(\|\hat{x}_k - x_k\|^2)} = \liminf_{k \rightarrow \infty} \frac{f_i(x_k) - f_i(\bar{x})}{\|\hat{x}_k - x_k\|^2}, \end{aligned}$$

contrary to (4). The proof is complete. □

**Remark 3.3.** The converse of Proposition 3.2 is not true in general. For example, let  $f: \mathbb{R} \rightarrow \mathbb{R}^2, x \mapsto (x^3, -x^3)$ ,  $g: \mathbb{R} \rightarrow \mathbb{R}, x \rightarrow -x^2$ , and  $C = -\mathbb{R}_+$ . Clearly,  $S = \mathbb{R}$  and  $\bar{x} = 0$  is an efficient solution of (MP). We claim that  $\bar{x}$  is not an essential local efficient solution of second-order. Indeed, if otherwise, there exist  $\gamma > 0$  and  $\delta > 0$  such that

$$\max\{f_1(x) - f_1(\bar{x}), f_2(x) - f_2(\bar{x}), \operatorname{dist}(g(x), C)\} \geq \gamma \|x - \bar{x}\|^2 \quad \forall x \in B_\delta(\bar{x}),$$

or, equivalently,  $\max\{x^3, -x^3\} \geq \gamma |x|^2 \quad \forall x \in (-\delta, \delta)$ . This implies that  $|x| \geq \gamma \quad \forall x \in (-\delta, \delta)$ , a contradiction.

The following result gives a sufficient optimality condition for an essential local efficient solution of second-order of problem (MP).

**Theorem 3.4.** *Let  $\bar{x}$  be a feasible solution of (MP). Suppose that for every  $u \in K(\bar{x}) \setminus \{0\}$  there exist  $\lambda \in \mathbb{R}_+^m$  and  $\mu \in \mathbb{R}^p$  not both zero such that the following conditions hold:*

$$\nabla_x L(\bar{x}, \lambda, \mu) = 0, \tag{6}$$

$$\nabla_{xx}^2 L(\bar{x}, \lambda, \mu)(u, u) + d^2 \delta_C(g(\bar{x}); \mu)(\nabla g(\bar{x})u) > 0. \tag{7}$$

Then  $\bar{x}$  is an essential local efficient solution of second-order of problem (MP).

*Proof.* The proof of the theorem follows some ideals of Benko et al. [18]. Suppose, for the sake of contradiction that  $\bar{x}$  is not an essential local efficient solution of second-order of problem (MP). Then

for each  $k \in \mathbb{N}$ , there exists  $x_k \in B_{1/k}(\bar{x})$  such that

$$\varphi(x_k) = \max\{f_1(x_k) - f_1(\bar{x}), \dots, f_m(x_k) - f_m(\bar{x}), \text{dist}(g(x_k), C)\} < \frac{1}{k} \|x_k - \bar{x}\|^2.$$

This implies that

$$f_i(x_k) - f_i(\bar{x}) \leq \frac{1}{k} \|x_k - \bar{x}\|^2, i = 1, \dots, m, \tag{8}$$

$$\text{dist}(g(x_k), C) \leq \frac{1}{k} \|x_k - \bar{x}\|^2. \tag{9}$$

Clearly,  $x_k \neq \bar{x}$  for all  $k \in \mathbb{N}$  and  $x_k \rightarrow \bar{x}$  as  $k \rightarrow \infty$ . For each  $k \in \mathbb{N}$ , put  $t_k := \|x_k - \bar{x}\|$  and  $u_k = t_k^{-1}(x_k - \bar{x})$ . Since  $\|u_k\| = 1$ , by passing a subsequence if necessary we may assume that  $u_k \rightarrow u$  with  $\|u\| = 1$  as  $k \rightarrow \infty$ . It follows from (8) that

$$\lim_{k \rightarrow \infty} \frac{f_i(x_k) - f_i(\bar{x})}{t_k} = \lim_{k \rightarrow \infty} \frac{\langle \nabla f_i(\bar{x}), t_k u_k \rangle + o(t_k)}{t_k} = \langle \nabla f_i(\bar{x}), u \rangle \leq 0 \tag{10}$$

and

$$\liminf_{k \rightarrow \infty} -\frac{f_i(\bar{x} + t_k u_k) - f_i(\bar{x})}{\frac{1}{2} t_k^2} \geq \lim_{k \rightarrow \infty} \frac{1}{k} = 0, i = 1, \dots, m. \tag{11}$$

By (9) and the closedness of  $C$ , for each  $k \in \mathbb{N}$ , there exists  $c_k \in C$  such that

$$\|c_k - g(x_k)\| = \text{dist}(g(x_k), C) \leq \frac{1}{k} t_k^2.$$

Put  $r_k := \frac{c_k - g(x_k)}{t_k^2}$  and  $v_k := \frac{g(x_k) - g(\bar{x}) + t_k^2 r_k}{t_k}$ . Then  $r_k \rightarrow 0$  as  $k \rightarrow \infty$  and

$$g(\bar{x}) + t_k v_k = g(x_k) + t_k^2 r_k = c_k \in C \text{ for all } k \in \mathbb{N}.$$

Moreover, by the differentiability of  $g$  one has

$$\lim_{k \rightarrow \infty} v_k = \lim_{k \rightarrow \infty} \frac{g(x_k) - g(\bar{x}) + t_k^2 r_k}{t_k} = \lim_{k \rightarrow \infty} \frac{t_k \langle \nabla g(\bar{x}), u_k \rangle + o(t_k) + t_k^2 r_k}{t_k} = \langle \nabla g(\bar{x}), u \rangle.$$

Hence  $\langle \nabla g(\bar{x}), u \rangle \in T_C(g(\bar{x}))$ . This and (10) imply that  $u \in K(\bar{x})$ . By the assumption of the theorem, there exist  $\lambda \in \mathbb{R}_+^m$  and  $\mu \in \mathbb{R}^p$  satisfying (6) and (7). By definition of the second subderivative and (11), we have

$$\begin{aligned} d^2 \delta_C(g(\bar{x}); \mu)(\nabla g(\bar{x})u) &= \liminf_{\substack{t \downarrow 0 \\ w \rightarrow \nabla g(\bar{x})u}} \frac{\delta_C(g(\bar{x}) + tw) - \delta_C(g(\bar{x})) - t \langle \mu, w \rangle}{\frac{1}{2} t^2} = \\ &= \liminf_{\substack{t \downarrow 0, w \rightarrow \nabla g(\bar{x})u \\ g(\bar{x}) + tw \in C}} \frac{-\langle \mu, w \rangle}{\frac{1}{2} t} \leq \liminf_{k \rightarrow \infty} \frac{-\langle \mu, t_k v_k \rangle}{\frac{1}{2} t_k^2} \\ &= \liminf_{k \rightarrow \infty} \frac{-\langle \mu, g(x_k) - g(\bar{x}) + t_k^2 r_k \rangle}{\frac{1}{2} t_k^2} \end{aligned}$$



$$\begin{aligned}
 &\leq \liminf_{k \rightarrow \infty} - \frac{\langle \lambda, f(x_k) - f(\bar{x}) \rangle}{\frac{1}{2} t_k^2} + \liminf_{k \rightarrow \infty} - \frac{\langle \mu, g(x_k) - g(\bar{x}) + t_k^2 r_k \rangle}{\frac{1}{2} t_k^2} \\
 &\leq \liminf_{k \rightarrow \infty} - \frac{L(x_k, \lambda, \mu) - L(\bar{x}, \lambda, \mu)}{\frac{1}{2} t_k^2} \\
 &= \liminf_{k \rightarrow \infty} - \frac{\nabla_x L(\bar{x}, \lambda, \mu)(t_k u_k) + \frac{1}{2} \nabla_{xx}^2 L(\bar{x}, \lambda, \mu)(t_k u_k, t_k u_k) + o(t_k^2)}{\frac{1}{2} t_k^2} \\
 &= -\nabla_{xx}^2 L(\bar{x}, \lambda, \mu)(u, u),
 \end{aligned}$$

contrary to (7). The proof is complete. □

The following result is a consequence of Theorem 3.4 and Proposition 3.2.

**Corollary 3.5.** *Let  $\bar{x}$  be a feasible solution of (MP). Suppose that for every  $u \in K(\bar{x})$  there exist  $\lambda \in \mathbb{R}_+^m$  and  $\mu \in \mathbb{R}^p$  not both zero such that conditions (6) and (7) hold. Then  $\bar{x}$  satisfies the quadratic growth condition.*

We finish this section by presenting an example to illustrate Theorem 3.4.

**Example 3.6.** Let  $f: \mathbb{R} \rightarrow \mathbb{R}^2, x \mapsto (x^2, -x^2)$ ,  $g: \mathbb{R} \rightarrow \mathbb{R}, x \rightarrow -x^2$ , and  $C = -\mathbb{R}_+$ . Then,  $S = g^{-1}(C) = \mathbb{R}$ . We now show that  $\bar{x} = 0$  is an essential local efficient solution of second-order of problem (MP). It is easy to check that  $K(\bar{x}) = \mathbb{R}$ . Choose  $\lambda = (1, 0)$  and  $\mu = 0$ . Then we have  $\nabla_x L(\bar{x}, \lambda, \mu) = 0$  and

$$\nabla_{xx}^2 L(\bar{x}, \lambda, \mu) + d^2 \delta_C(g(\bar{x}); \mu)(\nabla g(\bar{x})u) = \nabla_{xx}^2 L(\bar{x}, \lambda, \mu) + d^2 \delta_C(0; 0)(0) = 2 > 0.$$

Hence, by Theorem 3.4,  $\bar{x}$  is an essential local efficient solution of second-order of problem (MP).

#### 4. Conclusions

In this paper, we have presented a second-order sufficient optimality condition for an essential local efficient solution of second-order to nonconvex multiobjective optimization problems with operator constraint. It is meaningful if we can establish a necessary optimality condition for this problem that has no-gap between the proposed sufficient optimality condition. We aim to investigate this problem in future work.

#### References

- [1] J. F. Bonnans and A. Shapiro, *Perturbation analysis of optimization problems* (Springer series in operations research and financial engineering). New York, USA: Springer, 2000. doi: 10.1007/978-1-4612-1394-9.
- [2] R. Cominetti, “Metric regularity, tangent sets, and second-order optimality conditions”, *Appl. Math. Optim.*, vol. 21, pp. 265–287, Jan. 1990, doi: 10.1007/bf01445166.
- [3] N. Q. Huy and N. V. Tuyen, “New second-order optimality conditions for a class of differentiable optimization problems”, *J. Optim. Theory Appl.*, vol. 171, pp. 27–44, Jul. 2016, doi: 10.1007/s10957-016-0980-4.
- [4] N. Q. Huy, D. S. Kim, and N. V. Tuyen, “New second-order Karush–Kuhn–Tucker optimality conditions for vector optimization”, *Appl. Math. Optim.* vol. 79, pp. 279–307, Apr. 2019, doi: 10.1007/s00245-017-9432-2.

- [5] N. Q. Huy, B. T. Kien, G. M. Lee, and N. V. Tuyen, “Second-order optimality conditions for multiobjective optimization problems with constraints”, *Linear and Nonlinear Analysis*, vol. 5, no. 2, pp. 237–253, Sep. 2019.
- [6] N. H. Hung, H. N. Tuan, and N. V. Tuyen, “On second-order sufficient optimality conditions”, *HPU2. J. Sci.*, vol. 74, pp. 3–11, Aug. 2021.
- [7] N. H. Hung, H. N. Tuan, and N. V. Tuyen, “On the tangent sets of constraint systems”, *HPU2. Nat. Sci. Tech.*, vol. 1, no. 1, pp. 31–39, Aug. 2022, doi: 10.56764/hpu2.jos.2022.1.1.31-39.
- [8] A. Jourani, “Regularity and strong sufficient optimality conditions in differentiable optimization problems”, *Numer. Funct. Anal. Optim.*, vol. 14, no. 1–2, pp. 69–87, Jan. 1993, doi: 10.1080/01630569308816508.
- [9] D. S. Kim and N. V. Tuyen, “A note on second-order Karush-Kuhn-Tucker necessary optimality conditions for smooth vector optimization problems”, *RAIRO - Oper. Res.*, vol. 52, no. 2, pp. 567–575, Jul. 2018, doi: 10.1051/ro/2017026.
- [10] A. Mohammadi, B. S. Mordukhovich, and M. E. Sarabi, “Parabolic regularity in geometric variational analysis”, *Trans. Amer. Math. Soc.*, vol. 374, no. 3, pp. 1711–1763, Aug. 2021, doi: 10.1090/tran/8253.
- [11] J. P. Penot, “Second-order conditions for optimization problems with constraints”, *SIAM J. Control Optim.*, vol. 37, no. 1, pp. 303–318, Jan. 1998, doi: 10.1137/S0363012996311095.
- [12] R. T. Rockafellar and R. J. -B. Wets, *Variational analysis* (Grundlehren der mathematischen Wissenschaften). Heidelberg, Germany: Springer Berlin, 1998. doi: 10.1007/978-3-642-02431-3.
- [13] N. T. Toan, L. Q. Thuy, N. V. Tuyen, and Y. -B. Xiao, “Second-order KKT optimality conditions for multiobjective discrete optimal control problems”, *J. Global Optim.*, vol. 79, no. 1, pp. 203–231, Jan. 2021, doi: 10.1007/s10898-020-00935-7.
- [14] N. V. Tuyen, N. Q. Huy, and D. S. Kim, “Strong second-order Karush-Kuhn-Tucker optimality conditions for vector optimization”, *Appl. Anal.*, vol. 99, no. 1, pp. 103–120, Jun. 2018, doi: 10.1080/00036811.2018.1489956.
- [15] N. V. Tuyen, C. F. Wen, Y. B. Xiao, and J. C. Yao, “On second-order sufficient optimality conditions for  $C^1$  vector optimization problems”, *J. Nonlinear Convex Anal.*, vol. 23, no. 12, pp. 2859–2874, Dec. 2022.
- [16] Y. B. Xiao, N. V. Tuyen, J. C. Yao, and C. F. Wen, “Locally Lipschitz vector optimization problems: Second-order constraint qualifications, regularity condition, and KKT necessary optimality conditions”, *Positivity*, vol. 24, no. 2, pp. 313–337, Apr. 2020, doi: 10.1007/s11117-019-00679-z.
- [17] M. Benko and P. Mehlitz, “Why second-order sufficient conditions are, in a way, easy - or - revisiting calculus for second subderivatives”, *J. Convex Anal.*, vol. 30, no. 2, pp. 541–589, Jun. 2023.
- [18] M. Benko, H. Gfrerer, J. J. Ye, J. Zhang, and J. Zhou, “Second-order optimality conditions for general nonconvex optimization problems and variational analysis of disjunctive systems”, *SIAM J. Optim.*, vol. 33, no. 4, pp. 2625–2653, Oct. 2023, doi: 10.1137/22m1484742.
- [19] N. T. V. Hang and M. E. Sarabi, “A chain rule for strict twice epi-differentiability and its applications”, *SIAM J. Optim.*, vol. 34, no. 1, pp. 918–945, Feb. 2024, doi: 10.1137/22M1520025.
- [20] A. Mohammadi and M. E. Sarabi, “Twice epi-differentiability of extended-real-valued functions with applications in composite optimization”, *SIAM J. Optim.* vol. 30, no. 3, pp. 2379–2409, Jan. 2020, doi: 10.1137/19M1300066.